

Pediatric Dosage Chart

Tylenol (Acetaminophen)	Tylenol should be given every 4-6 hours. Do not give more than every 4 hours								
	6-11lbs	12-17lbs	18-23lbs	24-35lbs	36-47lbs	48-59lbs	60-71lbs	72-95lbs	95lbs <
Baby drops 80mg/0.8ml	1/2 dropper 0.4 ml	1 dropper	1 1/2 droppers 1.2 ml	2 droppers 1.6 ml	2 1/2 droppers 2.4 ml	N/A	N/A	N/A	N/A
Elixir or Syrup 160mg/5ml	1.25 ml	2.5 ml	4 ml	1 tsp	1.5 tsp	2 tsp	2.5 tsp	3 tsp	4 tsp
Chewable Tablets 80mg	N/A	N/A	1.5 tabs	2 tabs	3 tabs	4 tabs	5 tabs	6 tabs	8 tabs
Chewable Tablets 160mg	N/A	N/A	N/A	1 tab	1.5 tabs	2 tabs	2.5 tabs	3 tabs	4 tabs
Adult Tablets 325mg	N/A	N/A	N/A	0.5 tab	0.75 tab	1 tab	1.25 tab	1.5 tabs	2 tabs
Motrin/Advil (Ibuprofen)	Do not give Ibuprofen to children under 4 months of age. Ibuprofen should be given every 4-6 hours. Do not give doses more often than every 6 hours.								
	>11 lbs	12-17lbs	18-23lbs	24-35lbs	36-47lbs	48-59lbs	60-71lbs	72-95lbs	95lbs <
Infant Drops 50mg/1.25ml	N/A	1.25 ml	1.875 ml	2.5 ml	3.125 ml	3.725 ml	N/A	N/A	N/A
Suspension 100mg/5ml	N/A	2.5 ml (1/2 tsp)	4ml	1 tsp	1.5 tsp	2 tsp	2.75 tsp	3.25 tsp	4 tsp
Chewable Tablets 50mg	N/A	1 tab	1.5 tabs	2 tabs	3 tabs	4 tabs	5.5 tabs	6.5 tabs	8 tabs
Chewable Tablets 100mg	N/A	0.5 tab	1 tab	1 tab	1.5 tabs	2 tabs	2.5 tabs	3 tabs	4 tabs
Caplets Jr. 100mg	N/A	0.5 tab	1 tab	1 tab	1.5 tabs	2 tabs	2.5 tabs	3 tabs	4 tabs
Adult Tablets 200mg	N/A	N/A	N/A	0.5 tab	0.75 tab	1 tab	1 tab	1.5 tabs	2 tabs
Benadryl	Benadryl can cause children to be sleepy or sometimes more hyper than usual. Do not give more often than every 6 hours. Benadryl may not be very beneficial to young infants with cold symptoms.								
	>7lbs	7-13lbs	14-20lbs	21-27lbs	28-41lbs	42-55lbs	56-83lbs	84-111lbs	112lbs <
Syrup 12.5mg/5ml	N/A	2 ml	3 ml	4 ml	1 tsp	2 tsp	2.5 tsp	N/A	N/A
Chewable Tablets 12.5 mg	N/A	N/A	N/A	N/A	1 tab	2 tabs	2.5 tabs	3.5 tabs	N/A
Adult Caps	N/A	N/A	N/A	N/A	N/A	1 cap	1 cap	2 cap	2.5 cap

Note: 1 tsp = 5cc or 5ml 1/2 tsp = 2.5cc or 2.5 ml